

Globodera pallida	0	0	0	0	0	0	-	-	-	-	-	0	0	0	0	0	
Synchytrium endobioticum	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Clavibacter michiganensis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Potato spindle tuber viroid	0	0	0	0	0	0	-	-	-	-	-	0	0	0	0	0	
Tomato stolbur	0	0	0	0	0	0	-	-	-	-	-	-	-	-	-	-	
Meloidogyne chitwoodi and fallax	0	0	0	0	0	0	-	-	-	-	-	0	0	0	0	0	
Ditylenchus destructor	0	0	0	0	0	0	-	-	-	-	-	0	0	0	0	0	
3. Direct progeny tolerances																	
Other varieties and off types	0	0.01	0.25	0.25	0.5	0.5	0.25 & 0.1	0.25 & 0.1	0.25 & 0.1	0.25 & 0.1	0.5 & 0.2	0	0.01	0.1var 0.25 off	0.1 var 0.25 off	0.1 var 0.25 off	
Virus (%)	0	0.5	2 (1 severe)	4 (2 severe)	10 (5 severe)	10	4	4	4	4	10	0	0.05	4	4	4	

¹ Harmful organisms in bold are covered by the EU legislation on quarantine pests (Directive 2000/29/EC)

* The figure in brackets is the allowable % surface area covered: a tuber is deemed to be affected by the disease only if surface area affected exceeds the specified allowable surface tolerance.