


Quality control for drinking water production:

Monitoring and early warning



Dr Peter G Stoks RIWA Nieuwegein, NL

The Rhine

- 1320 km
- 185.000 km²
- >50 million people
- 9 riparian states
- multiple major industrial sites
- very dense shipping traffic





Around 25 million people depend on the Rhine for their drinking water (>10% in NL)

- Largely bank filtration as 1st treatment step
- Direct abstraction by 4 water works (2 in NL)



Knowledge about water quality important

- Sufficient for drinking water production?
- Long term → Monitoring program
- Short term → Early warning





- Aim:
 - Long term development (Trend)
 - Compliance testing
- Quality of data much more important than rapid availability
 - Periodical sampling
 - What happens inbetween?
 - Limited range of variables
 - What happens in addition?

Sudden spills not readily detected

→ Not suitable for operational purposes



Important catastrophies on the Rhine during the past decades

Mostly industrial

- Endosulfan (1968)
- Chloronitrobenzene (1970)
- Sandoz (1986)
- Numerous smaller incidents

Agricultural run-off

Isoproturon



ICPR Warning stations along

The Rhine

Mostly Water police stations

- "24/7 operation"
- Hotline to water authorities
- Info transmittal by standardized Fax forms





ICPR Warning stations along the Rhine

 Site-specific threshold values agreed upon

- Wide range of substances
- Eco-relevance and DW relevance





Important:

Warning stations can only transmit reported spills

- Industrial accidents
- Shipping accidents
- On-site measurements / observations

Not-reported events can only be detected by measurements

- Agricultural run-off
- Deliberate (illegal) spills
- Unnoticed accidents



Therefore: continuous Measurement stations along the Rhine

Mostly governmental stations, but also stations operated by water works

Examples: Weil am Rhein, Worms, Bimmen/Lobith, Nieuwegein



Early warning

- Aim:
 - Timely warnung (so as to allow measures)
- Rapid availability much more important than accuracy
 - continuous / high-frequency sampling
 - Very wide range of substances



Bimmen/Lobith facilities

(jointly operated by NL / DE authorities)

- General variables (pH, O₂, T, turbidity, conductivity....)
- Non-polar pollutants (GC/MS-Screening)
- Polar pollutants (LC/LCMS-Screening)
- Volatiles (P&T-GC/MS)
- BEWS (Daphnids, Algae)

Alarm thresholds set for drinking water production







Alarmeringswaarden 2011 voor het Nederlandse stroomgebied van Rijn en Maas

Parameter	Eenheld	Maas	Rijn
Algemeen	100	12	
 Zuurstof 	mg/l	<2	<2
 Zuurgraad 	рН	<6 of >9	<6 of >9
 Geleidendheid 	mS/m	100	3
 Troebelheid 	FTU	50	
Radioactiviteit			
 γ (per energie gebied) 	Bq/I	100	100
Zouten	- 1	NO SECTION	1704.0024
Chloride	mg/l	150	200
Fluoride	mg/l	1.5	12
Ammonium	mg N/I	4	:=
Metalen			
Cd	идл	3	
Cu	ид/1	15	
Pb Zn	µg/1 Г/дц	15 65	
Organische	PW.	164	
Microverontreinigingen			
Vluchtige verbindingen (SIVEVOC),	100510-000	1220	12.27
bekenden	μg/l	10	10
 Onbekenden 	μg/1	10	10
Diisopropylether	μg/l	20	
 A-polaire verbindingen (SIVEGOM), bekenden 	μg/I	3	3
 Onbekenden 	µg/1	3	3
 Tributylfosfaat 	μg/1	5	5
 Bestrijdingsmiddelen en afbraakproducten 	μg/l	1	0,3
Polaire verbindingen (SAMOS)	μg/I	3 (6 voor som) ¹	3
 Bestrijdingsmiddelen en afbraakproducten 	µg/1	1	0,3
Onbekenden	μg/I	3	3
Blosystemen			
 Algenmonitor 	% 2	4	4
Daphniasysteem	Tox index ³	10	10
Informatiegrenzen			
Onder kantooruren via e-mail adres			
Gamma straling	Bq/I	25	25
Bestrijdingsmiddelen	μg/l	0,5	0,1
Overige Stoffen (organisch)	μg/I	1	1
Chloride	mg/l		150

¹ de drinkwatersector wordt gewaarschuwd als de som van alle SAMOS componenten in het Lateraal kanaal boven de 6 ug/l utforming in de fluoresentie van de algen gedurende 2 uur groter of gelijk aan 4%. 2 percentage remming in de fluoresentie van de algen gedurende 2 uur groter of gelijk aan 4%. 3 gedurende 2 maal 2 minuten is de toxiciteitsindex in beide cuvetten groter of gelijk aan 10.

Postbus 17 E200 AA Leiysted Zuidenwagenplein 2 8224 AD Lelystad

Serekbaar vanaf station NG met budigs 3, 14, 143, 156 en interferer 315




NL overview

- Every 4th Dutchman depends on the Rhine for his drinking water
- 4 Water works supply this water
- 2 apply bank filtration
- 2 apply direct abstraction
- Direct abstraction very sensitive to pollution



Intake Amsterdam





Characteristics of a spill

- industrial spill / shipping accident: rapid concentration increase, short duration (hours / days)
- diffuse pollution (run-off): slow increase in conc, longer duration (weeks, months)




Direct abstraction: possible measures

- "dilution" with groundwater
- selective intake
- Interruption of intake



selective intake Nieuwegein





Intake stops Nieuwegein

Year	Number	Days	(Max)	Cause
1986	3	19	(9)	"Sandoz", 2,4-D, Acids
1987	1	3		Neopentylglycol
1988	3	20	(11)	Isophoron, MCPP, DCP
1989	1	4		Nitrobenzene
1990	1	6		Metamitron
1991-1993	0			
1994	1	36		Isoproturon
1995-1997	0			
1998	1	7		Isoproturon
1999	1	7		Isoproturon
2000	0			
2001	1	35		Isoproturon/Chlortoluron
2002	1	20		Isoproturon/Chlortoluron
2004	1	5		MTBE
2008	1	2		Dichlorobenzene



What is currently OK?

- Preventive action (governmental surveillance)
 - "Big Brother is watching you"
- Swift transferral of info
- Reliability / quality of information
 - Flow-time models, scope of analysis



What should improve?

- Finding causes for not-reported events
- Punishment / (financial) compensation
- Logistics (Fax, Email)
- References / Info about properties
- Research on better applicability of BEWS



adequate warning

more reliability in drinking water production